

PART III: CONSUMER INFORMATION

Fr Zoladex[®] LA goserelin depot

This leaflet is part III of a three-part "Product Monograph" published when ZOLADEX[®] LA was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about ZOLADEX LA. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What ZOLADEX LA is used for:

Prostate Cancer

ZOLADEX LA is used:

- For the palliative treatment of patients with hormone-dependent advanced carcinoma of the prostate (Stage D2).
- In combination with a non-steroidal antiandrogen and radiation therapy for the management of locally advanced (T3, T4) or bulky Stage T2b, T2c carcinoma of the prostate.
- As adjuvant hormone therapy to external beam irradiation for patients with locally advanced prostate cancer (Stage T3-T4).

Benign Conditions

ZOLADEX LA is indicated for the hormonal management of endometriosis, including pain relief and reduction of endometriotic lesions. Experience with ZOLADEX for the management of endometriosis has been limited to women 18 years of age and older, treated for 6 months.

What ZOLADEX LA does:

ZOLADEX LA treatment, given as recommended, results in suppression of your sex hormones (testosterone in men and estradiol in women). Any complaints you experience may be related to this hormone-suppressing action of ZOLADEX LA. These complaints may include hot flushes, swollen or tender breasts and reduction in sex drive.

When ZOLADEX LA should not be used:

You should not use ZOLADEX LA if:

- You are allergic to goserelin acetate or any nonmedicinal ingredients in ZOLADEX LA.
- You are a woman who has abnormal vaginal bleeding for an unknown reason.
- You are a woman who is pregnant.
- You are a woman who is breastfeeding.

What the medicinal ingredient is:

goserelin acetate

What the important nonmedicinal ingredients are:

Lactide-glycolide copolymer

What dosage forms ZOLADEX comes in:

ZOLADEX LA comes in a hard, cream-coloured, rod-shaped depot which contains 10.8mg of goserelin as goserelin acetate.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

ZOLADEX LA should be prescribed and managed by a doctor experienced with using this type of drug.

ZOLADEX LA may cause:

- Worsening of symptoms of prostate cancer at the beginning of the treatment (risk of spinal cord compression, or increased difficulty in urinating)
- Bone thinning (osteoporosis)

If you go into hospital, let the medical staff know you are receiving ZOLADEX LA.

In women, there are no clinical data on the effect of treating endometriosis with ZOLADEX LA for the periods in excess of six months.

ZOLADEX LA is not recommended for use in children.

ZOLADEX LA is unlikely to affect your ability to drive a car or to operate machinery.

Before you use ZOLADEX LA, talk to your doctor or pharmacist if any of the following applies to you:

- Have or have had any problem passing urine.
- Family history of severe osteoporosis (thinning of the bones with fractures).
- Have low bone mineral density (BMD).
- Taking other medicines that cause thinning of the bones.
- Have a low red blood cell count (anemia)
- Have heart disease, or have a heart condition called 'long QT syndrome', or a family history of this heart condition. ZOLADEX LA may increase the risk of having an abnormal heart rhythm (QT prolongation).
- Have diabetes.
- Are pregnant or planning to become pregnant. ZOLADEX LA should not be used during pregnancy, therefore, effective non-hormonal contraceptive methods should be used to prevent pregnancy during the treatment and until the return of menses after the last injection with ZOLADEX LA. After stopping ZOLADEX LA it may

take longer for some women to experience menses. Rarely, some women may enter menopause. If 12 weeks have passed after the last ZOLADEX LA injection and you do not experience menses, talk to your doctor.

INTERACTIONS WITH THIS MEDICATION

Check with your doctor or pharmacist before taking any other drugs, including non-prescription drugs (for colds, nausea, etc.) especially if you are taking any medicines for abnormal heart rhythm (antiarrhythmics) or medicines that may increase the risk of having an abnormal heart rhythm.

PROPER USE OF THIS MEDICATION

Usual Dose

- ZOLADEX LA is given as an injection under the skin of the abdomen by a trained health care professional, such as a doctor or nurse.
- **Prostate cancer:** one injection every 3 months
- **Endometriosis:** one injection every 12 weeks
- It is very important your doctor checks your progress at regular medical visits. Consult your doctor before you decide to change your treatment.
- If you need more information, ask your doctor.

Overdose

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose

If you missed your scheduled dose, contact your doctor for advice.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

As with all medicines, side effects are sometimes experienced with ZOLADEX LA.

Contact your doctor or pharmacist if you experience any of these problems:

- Tingling in your fingers or toes.
- Psychiatric problems such as hallucinations, disordered thoughts or personality change. These have occasionally been reported.
- There have been occasional reports of side effects with pituitary tumours. You may develop a tumour of the pituitary gland in your head or, if you have an existing tumour of the pituitary gland, ZOLADEX may cause it to bleed or collapse. Pituitary tumours may cause headaches, vomiting, loss of eyesight and unconsciousness.

- A local skin reaction may occur at the injection site such as pain, bruising, bleeding, itching, redness, burning and swelling. These reactions generally are mild and disappear after a few days. If they get worse or do not go away, tell your doctor.
- **Cancer patients:** Contact your doctor immediately if you develop: severe increased pain, numbness or weakness of the limbs, or persistent difficulty in urinating.

Use of ZOLADEX LA In Men

- When you first start receiving ZOLADEX LA you may feel some pain in your bones. If this happens tell your doctor and you may be given something for this.
- Very occasionally you may have trouble passing urine or experience lower back pain. If this happens, tell your doctor and you may be given something for this.
- You may experience hair loss, particularly the loss of body hair.

Use of ZOLADEX LA in Women

- For pre-menopausal women: menstruation stops with the 12 week depot of ZOLADEX LA. If regular menstruation persists, notify your doctor. Occasionally some women may enter menopause early, so when ZOLADEX LA treatment is stopped menstruation will not start again.
- Vaginal bleeding may occur. At the beginning of treatment, if you have fibroids a slight increase in symptoms, such as pain, may occur. These effects are usually short-lived and discontinue on continuation of treatment. If symptoms persist or you are uncomfortable, contact your doctor.
- ZOLADEX has been associated with the formation of ovarian cysts, which may cause pain for some women.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	

USE OF ZOLADEX LA IN MEN

Very Common (more than 10 in every 100 patients are likely to have them)

Reduced sex drive/impotence	√		
Hot flushes and sweating	√		

Common (1 to 10 in every 100 patients are likely to have them)

Change in breast size	√		
Injection site reaction	√		
Bone pain	√		
Rises in blood sugar levels		√	
Tingling in fingers or toes	√		
Changes in blood pressure		√	
Skin rashes	√		
Thinning of bones		√	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Heart failure (reduced heart function) or heart attack		√	
Weight gain	√		
Uncommon (1 to 10 in every 1000 patients are likely to have them)			
Tender breasts	√		
Joint pain		√	
Allergic reactions		√	
USE OF ZOLADEX LA IN WOMEN			
Very Common (more than 10 in every 100 patients are likely to have them)			
Reduced sex drive	√		
Hot flushes and sweating	√		
Vaginal dryness	√		
Change in breast size	√		
Injection site reaction	√		
Common (1 to 10 in every 100 patients are likely to have them)			
Mood changes including depression		√	
Tingling in fingers and toes	√		
Headache		√	
Changes in blood pressure		√	
Skin rashes	√		
Thinning of bones		√	
Joint pain		√	
Hair loss (usually mild but occasionally severe)	√		
Weight gain	√		
Uncommon (1 to 10 in every 1000 patients are likely to have them)			
Allergic reactions		√	

This is not a complete list of side effects. For any unexpected effects while taking ZOLADEX[®] LA, contact your doctor or pharmacist.

HOW TO STORE IT

- ZOLADEX LA should not be used after the expiry date on the pack. Store ZOLADEX LA in its original pack between 2°C and 25°C.
- If your doctor decides to stop your treatment, return ZOLADEX LA to the pharmacy for proper disposal
- Keep your ZOLADEX LA in a safe place where children cannot reach it. It could harm them.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701D
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

NOTE: This CONSUMER INFORMATION leaflet provides you with the most current information at the time of printing.

For the most current information, the Consumer Information Leaflet plus the full Product Monograph, prepared for health professionals can be found at:

www.astrazeneca.ca,
or by contacting the sponsor, AstraZeneca Canada Inc. at:
Customer Inquiries – 1 (800) 668-6000,
Renseignements – 1 (800) 461-3787.

This leaflet was prepared by:
AstraZeneca Canada Inc. Mississauga, Ontario L4Y 1M4

ZOLADEX[®] and the AstraZeneca logo are registered trademarks of the AstraZeneca group of companies.

© AstraZeneca 2000, 2009, 2010

Last revised: March 4, 2011